



## Rappels de trigonométrie

### Valeurs remarquables

$x$	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$
$\cos x$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
$\sin x$	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
$\tan x$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	$\infty$
$\cotan x$	$\infty$	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0

### Symétries des fonctions trigonométriques

$\cos(-x) = \cos(x)$	$\sin(-x) = -\sin(x)$	$\tan(-x) = -\tan(x)$
$\cos(\pi + x) = -\cos(x)$	$\sin(\pi + x) = -\sin(x)$	$\tan(\pi + x) = \tan(x)$
$\cos(\pi - x) = -\cos(x)$	$\sin(\pi - x) = \sin(x)$	$\tan(\pi - x) = -\tan(x)$
$\cos\left(\frac{\pi}{2} - x\right) = \sin(x)$	$\sin\left(\frac{\pi}{2} - x\right) = \cos(x)$	$\tan\left(\frac{\pi}{2} - x\right) = \cotan(x)$
$\cos\left(\frac{\pi}{2} + x\right) = -\sin(x)$	$\sin\left(\frac{\pi}{2} + x\right) = \cos(x)$	$\tan\left(\frac{\pi}{2} + x\right) = -\cotan(x)$

### Formules d'addition

<b>formules d'addition</b>	$\cos(a + b) = \cos a \cos b - \sin a \sin b$	$\sin(a + b) = \sin a \cos b + \cos a \sin b$	$\tan(a + b) = \frac{\tan a + \tan b}{1 - \tan a \tan b}$
	$\cos(a - b) = \cos a \cos b + \sin a \sin b$	$\sin(a - b) = \sin a \cos b - \cos a \sin b$	$\tan(a - b) = \frac{\tan a - \tan b}{1 + \tan a \tan b}$
<b>cas des arcs doubles</b>	$\cos 2a = 2\cos^2 a - 1 = 1 - 2\sin^2 a$ $= \cos^2 a - \sin^2 a$	$\sin 2a = 2\cos a \sin a$	$\tan 2a = \frac{2\tan a}{1 - \tan^2 a}$
<b>expressions en fonction de <math>t = \tan \frac{a}{2}</math></b>	$\cos a = \frac{1 - t^2}{1 + t^2}$	$\sin a = \frac{2t}{1 + t^2}$	$\tan a = \frac{2t}{1 - t^2}$

### Conséquences des formules d'addition :

$$\begin{aligned}\cos a \cos b &= \frac{1}{2}(\cos(a + b) + \cos(a - b)) \\ \sin a \sin b &= \frac{1}{2}(\cos(a - b) - \cos(a + b)) \\ \sin a \cos b &= \frac{1}{2}(\sin(a + b) + \sin(a - b))\end{aligned}$$

### Équations trigonométriques

$$\cos a = \cos b \Leftrightarrow \begin{cases} a \equiv b [2\pi] \\ \text{ou} \\ a \equiv -b [2\pi] \end{cases}$$

$$\sin a = \sin b \Leftrightarrow \begin{cases} a \equiv b [2\pi] \\ \text{ou} \\ a \equiv \pi - b [2\pi] \end{cases}$$

$$\tan a = \tan b \Leftrightarrow a \equiv b [\pi]$$